

References

- [1] Ge PS, Runyon BA. The changing role of beta-blocker therapy in patients with cirrhosis. *J Hepatol* 2014;60:643–653.
- [2] Lebre D, Nouel O, Corbic M, Benhamou JP. Propranolol – a medical treatment for portal hypertension? *Lancet* 1980;2:180–182.
- [3] Abrahams JG, Tarantino I, Turnes J, Garcia-Pagan JC, Rodes J, Bosch J. Hemodynamic response to pharmacological treatment of portal hypertension and long-term prognosis of cirrhosis. *Hepatology* 2003;37:902–908.
- [4] Hernandez-Gea V, Aracil C, Colomo A, Garupera I, Poca M, Torras X, et al. Development of ascites in compensated cirrhosis with severe portal hypertension treated with beta-blockers. *Am J Gastroenterol* 2012;107:418–427.
- [5] Thalheimer U, Bosch J, Burroughs AK. How to prevent varices from bleeding: shades of grey – the case for nonselective beta blockers. *Gastroenterology* 2007;133:2029–2036.
- [6] Thalheimer U, Triantos CK, Samonakis DN, Patch D, Burroughs AK. Infection, coagulation, and variceal bleeding in cirrhosis. *Gut* 2005;54:556–563.
- [7] Farland MZ, Peters CJ, Williams JD, Bielak KM, Heidel RE, Ray SM. Beta-Blocker use and incidence of chronic obstructive pulmonary disease exacerbations. *Ann Pharmacother* 2013;47:651–656.
- [8] Short PM, Lipworth SL, Elder DH, Schembri S, Lipworth BJ. Effect of beta blockers in treatment of chronic obstructive pulmonary disease: a retrospective cohort study. *BMJ* 2011;342:d2549.
- [9] Paravastu SC, Mendonca DA, Da SA. Beta blockers for peripheral arterial disease. *Cochrane Database Syst Rev* 2013;9:CD005508.
- [10] Ryden L, Standl E, Bartnik M, Van den Berghe G, Betteridge J, de Boer MJ, et al. Guidelines on diabetes, pre-diabetes, and cardiovascular diseases: executive summary. The Task Force on Diabetes and Cardiovascular Diseases of the European Society of Cardiology (ESC) and of the European Association for the Study of Diabetes (EASD). *Eur Heart J* 2007;28:88–136.

- [11] Singh V, Singh A, Singh B, Vijayvergiya R, Sharma N, Ghai A, et al. Midodrine and clonidine in patients with cirrhosis and refractory or recurrent ascites: a randomized pilot study. *Am J Gastroenterol* 2013;108:560–567.
- [12] Triantos C, Samonakis D, Thalheimer U, Patch D, Burroughs A. The relationship between liver function and portal pressure: what comes first, the chicken or the egg? *J Hepatol* 2005;42:146–147.

Ulrich Thalheimer*

The Exeter Liver Unit,

Royal Devon & Exeter Foundation Trust,

Exeter, UK

*Corresponding author.

E-mail address: uthalheimer@gmx.net

Jaime Bosch

Hepatic Hemodynamic Laboratory,

Hospital Clínic-IDIBAPS and Ciberehd,

University of Barcelona, C.Villarreal 170,

08036 Barcelona,

Spain

Andrew K. Burroughs

Sheila Sherlock Liver Centre,

Royal Free Hospital,

London, UK



Beta-blockers in cirrhosis: Thank you for your attention

To the Editor:

We graciously thank Drs. Ferrarese, Thalheimer, their colleagues, and the editorial board and worldwide readership of the *Journal of Hepatology* for their interest in our article [1]. We are humbled and honored by the international attention that our article has received, and are excited by the debate that it has ignited within the hepatology community. We have received numerous positive correspondences regarding our review [2]. In light of recent studies that have stirred controversy, and in anticipation of future studies that will continue to stir controversy, we believed a fresh objective look at the emerging evidence in the use of beta-blockers in cirrhosis was warranted.

At our institution, our chief executive officer frequently impresses his mantra to “put patients first” upon our entire hospital staff [3]. As physicians, we all believe that what comes first is our relentless and selfless service and dedication to our patients. Sometimes, this involves challenging existing treatments when they are later found to be harmful, such as when the same investigator who first studied the benefits of beta-blockers in patients with cirrhosis no longer found these benefits universally applicable [4].

We have frequently cared for patients with advanced cirrhosis who were seemingly harmed by beta-blockers once they had fallen outside a certain “therapeutic window” (that had only recently been hypothesized) [5]. Once their beta-blockers were discontinued, azotemia, hypotension, and acute kidney injury frequently and convincingly resolved. We were certain other

clinicians around the world must have encountered similar experiences, yet the scientific evidence appeared to be lacking. How could we turn our back on 30 years of highly cited landmark studies that promoted the use of beta-blockers? Did we miss something that was actually harming our patients?

Our article was therefore inspired by a need to re-explore the data. We do not dispute that the benefits of beta-blockers have been well-documented in patients with cirrhosis; however, that was not the focus of our article. We acknowledged these existing benefits and set out to more closely examine the studies that had been ignored [6], forgotten [7], or downplayed [8,9]. What we found was evidence – the quality of which can be debated, but nonetheless evidence stemming from astute clinicians making clinical observations – that beta-blockers were perhaps not as universally indicated as even we ourselves had previously believed. Just as the acetylsalicylic acid of cardiology has its limitations, the “aspirin of hepatology” appears to have its own pitfalls and limitations.

Even now, as this debate re-emerges, a new study from Mandorfer and colleagues provides fresh evidence demonstrating the detrimental effect of beta-blocker treatment after the development of spontaneous bacterial peritonitis [10]. More studies and debate are certain to follow.

We are extremely pleased that our review has achieved its intended effect of renewing dialogue and reopening the scrutiny on beta-blockers. We hope that this dialogue, along with new research specifically focusing at studying the end-stages of

Letters to the Editor

cirrhosis, will continue to be encouraged by the international hepatology community.

Conflict of interest

The authors declared that they do not have anything to disclose regarding funding or conflict of interest with respect to this manuscript.

References

- [1] Ge PS, Runyon BA. The changing role of beta-blocker therapy in patients with cirrhosis. *J Hepatol* 2014;60:643–653.
- [2] Zaman A. NEJM Journal Watch. Beta-Blocker Therapy in Cirrhosis: A Paradigm Shift. <http://www.jwatch.org/na33999/2014/03/18/beta-blocker-therapy-cirrhosis-paradigm-shift>. Accessed April 13, 2014.
- [3] UC Regents. CEO of hospital system has mantra: Put patients first. <http://www.today.ucla.edu/portal/ut/PRN-ceo-of-hospital-system-has-mantra-96848.aspx>. Accessed April 13, 2014.
- [4] Lebrec D, Poynard T, Hillon P, et al. Propranolol for prevention of recurrent gastrointestinal bleeding in patients with cirrhosis: a controlled study. *N Engl J Med* 1981;305:1371–1374.
- [5] Krag A, Wiest R, Albillos A, et al. The window hypothesis: haemodynamic and non-haemodynamic effects of beta-blockers improve survival of patients with cirrhosis during a window in the disease. *Gut* 2012;61:967–969.
- [6] Krag A, Bendtsen F, Henriksen JH, et al. Low cardiac output predicts development of hepatorenal syndrome and survival in patients with cirrhosis and ascites. *Gut* 2010;59:105–110.
- [7] Llach J, Gines P, Arroyo V, et al. Prognostic value of arterial pressure, endogenous vasoactive systems, and renal function in cirrhotic patients admitted to the hospital for the treatment of ascites. *Gastroenterology* 1988;94:482–487.
- [8] Serste T, Francoz C, Durand F, et al. Beta-blockers cause paracentesis-induced circulatory dysfunction in patients with cirrhosis and refractory ascites: a cross-over study. *J Hepatol* 2011;55:794–799.
- [9] Serste T, Melot C, Francoz C, et al. Deleterious effects of beta-blockers on survival in patients with cirrhosis and refractory ascites. *Hepatology* 2010;52:1017–1022.
- [10] Mandorfer M, Bota S, Schwabl P, et al. Non-selective beta blockers increase risk for hepatorenal syndrome and death in patients with cirrhosis and spontaneous bacterial peritonitis. *Gastroenterology* 2014, pii: S0016-5085(14)00306-0, <http://dx.doi.org/10.1053/j.gastro.2014.03.005>. [Epub ahead of print].

Phillip S. Ge
Bruce A. Runyon*

Division of Digestive Diseases, David Geffen School of Medicine at
UCLA, Los Angeles, CA, USA

*Corresponding author.

E-mail address: barunyon@mednet.ucla.edu



Is the pathway of energy metabolism modified in advanced cirrhosis?

To the Editor:

The role of metabolic alterations and adaptations is becoming increasingly evident in the pathogenesis of cirrhosis and hepatocellular carcinoma (HCC) [1–4]. Using a rat model of cirrhosis, Nishikawa *et al.* [5] demonstrate that early stage cirrhotic hepatocytes switch to glycolysis to meet their energy requirements as a result of a decline in oxidative phosphorylation, but that this mechanism fails in late cirrhosis. Since HCC typically arises in the background of liver cirrhosis [6] and also encompasses a glycolytic phenotype [7], it would be important to determine how this is related to the altered metabolism of cirrhotic hepatocytes. Since early cirrhotic cells are more glycolytic than advanced or failing cirrhotic cells, it remains unclear as to whether the progression of HCC is facilitated by metabolically active early cirrhotic cells or represents an escape mechanism of late cirrhotic hepatocytes which are unable to sustain their energy demands.

Conflict of interest

The authors declare that they do not have anything to disclose regarding funding or conflict of interest with respect to this manuscript.

References

- [1] Krahenbuhl S, Reichen J. Adaptation of mitochondrial metabolism in liver cirrhosis. Different strategies to maintain a vital function. *Scand J Gastroenterol Suppl* 1992;193:90–96.
- [2] Geschwind JF, Georgiades CS, Ko YH, Pedersen PL. Recently elucidated energy catabolism pathways provide opportunities for novel treatments in hepatocellular carcinoma. *Expert Rev Anticancer Ther* 2004;4:449–457.
- [3] Aram G, Potter JJ, Liu X, Wang L, Torbenson MS, Mezey E. Deficiency of nicotinamide adenine dinucleotide phosphate, reduced form oxidase enhances hepatocellular injury but attenuates fibrosis after chronic carbon tetrachloride administration. *Hepatology* 2009;49:911–919.
- [4] Ganapathy-Kanniappan S, Kunjithapatham R, Geschwind JF. Glyceraldehyde-3-phosphate dehydrogenase: a promising target for molecular therapy in hepatocellular carcinoma. *Oncotarget* 2012;3:940–953.
- [5] Nishikawa T, Bellance N, Damm A, Bing H, Zhu Z, Handa K, et al. A switch in the source of ATP production and a loss in capacity to perform glycolysis are hallmarks of hepatocyte failure in advanced liver disease. *J Hepatol* 2014;60:1203–1211.
- [6] Fattovich G, Stroffolini T, Zagni I, Donato F. Hepatocellular carcinoma in cirrhosis: incidence and risk factors. *Gastroenterology* 2004;127:S35–S50.
- [7] Beyoglu D, Idle JR. The metabolomic window into hepatobiliary disease. *J Hepatol* 2013;59:842–858.

Shanmugasundaram Ganapathy-Kanniappan*

Swathi Karthikeyan

Jean-Francois Geschwind

Russell H. Morgan Department of Radiology and Radiological
Sciences, The Johns Hopkins University School of Medicine,
Baltimore, MD, USA

*Corresponding author.

E-mail address: gshanmu1@jhmi.edu

Esteban Mezey

Department of Medicine, The Johns Hopkins
University School of Medicine, Baltimore, MD, USA